



Archived at the Flinders Academic Commons:

<http://dspace.flinders.edu.au/dspace/>

'This is the peer reviewed version of the following article:
Land, Gordon, & Watt. (2017). Isokinetic clinical assessment
of rotator cuff strength in subacromial shoulder
impingement. *Musculoskeletal Science and Practice*, 27,
32-39.

which has been published in final form at

<http://dx.doi.org/10.1016/j.msksp.2016.11.012>

© 2017 Elsevier. This manuscript version is made available
under the CC-BY-NC-ND 4.0 license [http://
creativecommons.org/licenses/by-nc-nd/4.0/](http://creativecommons.org/licenses/by-nc-nd/4.0/)

Background: Current conservative management of subacromial shoulder impingement (SSI) includes generic strengthening exercises, especially for internal (IR) and external (ER) shoulder rotators. However, there is no evidence that the strength or the ratio of strength between these muscle groups is different between those with SSI (cases) and an asymptomatic population (controls).

Objective: To identify if isokinetic rotator cuff strength or the ratio of strength is significantly different between cases and controls.

Study Design: Case Control Study.

Method: Fifty one cases with SSI and 51 asymptomatic controls matched for age, gender, hand dominance and physical activity level completed isokinetic peak torque glenohumeral IR and ER testing. Within the SSI group, 31 dominant limbs were symptomatic and 20 non-dominant limbs were symptomatic. IR and ER were measured separately using continuous reciprocal concentric (con) and eccentric (ecc) contraction cycles at a speed of 60⁰ degrees per second and again at 120⁰ degrees per second. Values of peak torque (PT), relative peak torque (RPT) and ratios were compared using independent t-tests between the SSI and asymptomatic groups.

Results: Significant strength differences between the two groups were present only when the symptomatic SSI shoulder was the dominant shoulder (con ER PT at 60⁰ /second, ecc ER PT at 120⁰ /second, ecc ER RPT at 120⁰ /second and ecc IR PT at 60⁰ /second and 120⁰ /second).

Conclusions: Changes in rotator cuff strength in SSI may be related to limb dominance, which may have implications for strengthening regimes.

Level of Evidence: Level 3a

Key Words: *Isokinetic, Peak Torque, Glenohumeral, Rotation, Impingement*

INTRODUCTION

Subacromial shoulder impingement (SSI) is a common condition characterized by anterolateral catching pain or aching of the shoulder, without a history of trauma. Pain originates from the tissues within the subacromial space including the rotator cuff (N. Hanchard, Cummins, & Jeffries, 2004; J. S. Lewis, Green, & Dekel, 2001). In people with SSI it is proposed rotator cuff muscle weakness develops secondary to inflammation and degeneration that occurs as a result of mechanical compression from a structure external to the tendon, known as extrinsic SSI (Michener, McClure, & Karduna, 2003), or as a result of overuse and tension overload affecting the tendons intrinsically, as in tendinopathy, known as intrinsic SSI (Jeremy S Lewis, 2009).

The rotator cuff musculature stabilise as well as move the glenohumeral joint. Subscapularis acts as an internal rotator and infraspinatus, teres minor and supraspinatus act as external rotators (Dark, Ginn, & Halaki, 2007; M.M. Reinold et al., 2004). The rotator cuff has been shown to produce different activity levels dependent on the direction of movement (J. Lewis & Ginn, 2015) and the rotator cuff and biceps have been identified to pre-set prior to actual movement being performed in asymptomatic young male shoulders (David et al., 2000).

Current conservative management of SSI includes generic resistance band strengthening exercises for the rotator cuff particularly shoulder external rotators (ER) (Holmgren, Bjornsson Hallgren, Oberg, Adolfsson, & Johansson, 2012; Kuhn, 2009). Exercises prescribed for SSI appear to be based on results from EMG studies and the experience and general knowledge of the physiotherapist (Dark et al., 2007; Holmgren et al., 2012; Michael.M. Reinold et al., 2007; Tate, McClure, Young,

Salvatori, & Michener, 2010). Previous isokinetic studies comparing rotator cuff strength in a diagnosed SSI group with an asymptomatic group analysed within group differences of the (1) painful versus non-painful shoulder in those with SSI and (2) dominant versus non-dominant shoulder in an asymptomatic group and then (3) compared the values from these two analyses (Leroux et al., 1994; MacDermid, Ramos, Drosdowech, Faber, & Patterson, 2004; Tyler, Nahow, Nicholas, & McHugh, 2005). Although comparison of dominant and non-dominant limbs have been reported there is no indication that matched dominance was considered in recruitment of symptomatic and asymptomatic group participants in these studies. Greater strength in the dominant upper limb compared to the non-dominant upper limb of the asymptomatic group is expected however this may or may not be the case in a SSI population. Lack of matching for arm dominance limits the opportunity to understand specific variations in strength which may be present due to usual physical activities. Matching of dominance should be an essential component to understand upper limb isokinetic testing results.

Isokinetic testing, performed through an active range at a constant velocity, is a reliable measure of shoulder strength (Land & Gordon, 2011). Internal and external rotation are consistently used to assess the rotator cuff, (Ludewig & Cook, 2000; Reddy, Mohr, Pink, & Jobe, 2000) with bilateral comparison of concentric peak torque shown to be the most appropriate outcome parameter for comparisons between healthy subjects and those with a painful condition (van Meeteren, Roebroek, Selles, Stijnen, & Stem, 2004). A seated testing position with the shoulder positioned in the scapular plane is reported to optimize the length tension relationship of the rotators, maximizing conformity between the humeral head and glenoid and is the most comfortable testing position (Kuhlman et al., 1992).

Functionally, EMG studies have identified that during internal rotation pectoralis major muscle activity is greater than subscapularis which is greater than latissimus dorsi expressed as a percentage of maximum voluntary isometric contraction (%MVIC) at low, medium and high exercise intensities (Dark et al., 2007). During external rotation infraspinatus, teres minor and supraspinatus muscle activity (%MVIC) is much greater than deltoid muscle activity at all exercise intensities and when the arm is positioned in the scapular plane (Dark et al., 2007; M.M. Reinold et al., 2004).

Strength changes in SSI not only result from decreased use of the shoulder to avoid pain but also due to altered motor strategies (Roy, Moffet, & McFadyen, 2008); decreased central motor corticospinal excitability when symptoms are ≥ 12 months (Ngomo, Mercier, Bouyer, Sacoie, & Roy, 2015); and inhibition when low to moderate pain levels are present (Dube & Mercier, 2011). Understanding possible muscle strength changes will assist treating clinicians to provide targeted exercise programs and enhance recovery.

The purpose of this study was to compare rotator cuff strength and strength ratios in a group diagnosed with SSI (cases) and a control group, matched for age, gender, hand dominance and physical activity level. The hypothesis was that there would be a difference in muscle strength between the painful shoulder in the SSI group and the dominance matched shoulder in the control group.

METHOD

A case control study, using matched pairs, was conducted to compare rotator cuff muscle strength in those with positive signs of SSI, of gradual onset and without

trauma, to an asymptomatic control matched for age, gender, hand dominance and physical activity level.

All testing was performed by an experienced musculoskeletal physiotherapist with over 20 years clinical experience, with both shoulders being measured in all participants.

The recruitment, inclusion and exclusion criteria for this case control study have been previously reported and are provided here for the convenience of the reader.

Participant Information and Consent

Ethical approval for this study was granted by the James Cook University (JCU) Human Ethics Committee (approval: H3945). Written informed consent was obtained from each of the participants.

Participants were recruited from the Townsville community and clients presenting to the JCU Physiotherapy Clinic between June 2011 and July 2013. Recruitment for both groups was via emails and word of mouth throughout the University staff, students and their extended networks. In addition, cases were also recruited using an advertisement in the local Townsville press and in the waiting area of the clinic. Cases identified with the advertisement 'Do you feel a sharp catch in your shoulder when raising your arm which eases when you lower your arm down? Is this making it difficult for you to wash your hair or reach up into an overhead cupboard or get your shirt on easily? Is it becoming painful to lie directly onto that shoulder at night?' They then contacted the investigator who arranged an assessment to determine eligibility. Controls were asked to be between 40 and 60 years of age with no history of shoulder, neck or upper back injuries and no reports of painful symptoms in any of

these areas in the previous twelve months. Both groups were required to meet the inclusion criteria.

Power Analysis

This study was part of a larger study in which a pre-study sample size calculation was performed, with $\alpha = 0.05$ and power 0.8, (Altman, 1991) which identified a minimum of 45 cases and 45 controls were needed. This sample size was adequate when compared with a calculation based on an isokinetic study comparing rotator cuff strength in a diagnosed SSI group with an asymptomatic group, peak torque external rotation at 60 degrees per second (mean difference 10Nm, standard deviation 2Nm (Leroux et al., 1994)) .

Inclusion and Exclusion Criteria

Forty to 60 year old participants were recruited to reflect the reported peak age for shoulder impingement (Ostor, Richards, Prevost, Speed, & Hazleman, 2005; van der Windt, Koes, de Jong, & Bouter, 1995). Symptom free volunteers as well as people with unilateral shoulder pain completed a screening questionnaire to determine their eligibility for this study. The questionnaire was used to exclude participants, in both the case and control groups, who had:

- Been participating in intense shoulder strength training during the 6 months prior to entering the study. This was defined as high load upper body weight training two or more times per week.
- Recent (within previous two years) or current pregnancy. This exclusion was necessary due to the effect of ligamentous laxity and postural changes associated with pregnancy.

- Previously undergone shoulder surgery or suffered a fracture of the shoulder girdle
- Glenohumeral instability identified by a grade 2 or 3 anterior, posterior or inferior load and shift test (assessed objectively) or a history of shoulder dislocation
- Scoliosis (also observed visually)
- Been experiencing cervical or thoracic pain currently or in the previous six months
- Diagnosed systemic or neurological disease (Type 2 diabetes was not screened for)
- Shoulder corticosteroid injection at any time in the past

If the questionnaire indicated they were eligible, a physical assessment was conducted of both the case and control volunteers.

In order to rule out other shoulder diagnoses and focus only on SSI, case group participants had:

- a minimum of three positive orthopaedic special tests (Michener, Walsworth, Doukas, & Murphy, 2009; Park, Yokota, Gill, Rassi, & McFarland, 2005). Hawkins-Kennedy (Hawkins & Kennedy, 1980) and/or Neer (Neer, 1983) must be positive along with two of the following: external rotation resistance test (Michener et al., 2009), tendon palpation (N. Hanchard et al., 2004), horizontal (cross-body) adduction (Park et al., 2005), painful arc (Kessel & Watson, 1977), drop arm test (Park et al., 2005), Yergason test (Dalton, 1989), Speed test (Dalton, 1989; Park et al., 2005)
- 'catching' or aching pain without appreciable joint stiffness (N. C. A. Hanchard & Handoll, 2008)

- a painful arc elicited with pain easing on lowering the arm (N. Hanchard et al., 2004)
- pain localized to the anterior or antero-lateral-superior shoulder (J. S. Lewis et al., 2001)
- insidious onset of symptoms with a possible history of gradual progression over time but without history of trauma (Bigliani & Levine, 1997)
- xray or ultrasound scans revealing osteophytes within the subacromial region, calcification of tendons or large rotator cuff tears . Alterations in acromial shape and bursal thickening were noted but did not prevent inclusion

Procedure

The shoulder pain and disability index (SPADI) was completed to further describe the SSI group. This outcome measures pain and disability associated with shoulder impairment (Roach, Budiman-Mak, Songsiridej, & Lertratanakul, 1991) and is frequently used for assessment of SSI syndrome (Dogu, Sahin, Ozmaden, Yilmaz, & Kuran, 2013). The visual analogue scale (VAS) was used to measure pain at rest and during activity (Jensen, Karoly, & Braver, 1986). Physical activity level was established by completing the short form of The International Physical Activity Questionnaire (IPAQ) (Craig et al., 2003). The IPAQ assesses three specific types of activity (1) walking (2) moderate-intensity activities such as cycling for transport and yard work (3) vigorous intensity activities such as running and boxing. A rating of low, medium or high physical activity is given for the duration (in minutes) and frequency (days) of activity.

Isokinetic testing was performed using a Humac Norm Computerised Dynamometer ((CSMI), 2006). Isokinetic reliability studies were completed prior to data collection. The testing method has been shown to be reliable when testing a group experiencing SSI and an asymptomatic group (MacDermid et al., 2004) and has been used in similar studies (Dulgeroglu, Kirbiyik, Ersoz, & Ozel, 2013; Erol, Ozcakar, & Celiker, 2008; Leroux et al., 1994; Tyler et al., 2005).

Isokinetic peak torque glenohumeral internal (IR) and external rotation (ER) were measured separately using continuous reciprocal concentric and eccentric contraction cycles at a speed of 60⁰ degrees per second and again at 120⁰ degrees per second. Testing was performed through a total range of 60 degrees from neutral rotation. Neutral rotation to 30⁰ IR and from neutral rotation to 30⁰ ER. Gravity correction was not applied as the range of motion tested in the seated position resulted in gravity affecting both IR and ER movements equally. Further, as significant error has been found when applying gravity correction due to the inability of the person to relax it was not considered advantageous (Bygott, McMeeken, Carroll, & Story, 2001). (Full details of method in Appendix A).

Data Analysis

Data were analysed using IBM SPSS Statistics Version 22. Descriptive statistics (mean, standard deviation, range) were calculated for each variable. All data was tested and found to be normally distributed.

A reliability study was analysed using intra-class correlation.

The measurements included in analyses were:

1. Peak torque of isokinetic concentric and eccentric ER and IR measured in Newton Metres

2. Relative peak torque of isokinetic concentric and eccentric ER and IR. This was calculated as peak torque divided by individual's body weight

3. Ratio of eccentric peak torque ER to concentric IR peak torque =

$$\frac{\text{eccentric peak torque ER}}{\text{concentric peak torque IR}}$$

4. Ratio of concentric peak torque ER peak torque to concentric IR peak torque=

$$\frac{\text{concentric peak torque ER}}{\text{concentric peak torque IR}}$$

Comparisons between matched SSI cases and controls were completed using independent samples t-tests, with significance $p \leq 0.05$. When the dominant shoulder was painful in the SSI group it was compared to the dominant shoulder in the control group and when the non-dominant shoulder was painful in the SSI group it was compared to the non-dominant shoulder in the control group.

RESULTS

An isokinetic reliability study completed on an asymptomatic group prior to data collection indicated high intra-rater reliability for all measures (ICC 0.948, CI 0.992 to 0.965) (Table 1).

TABLE 1: OUTCOME OF RELIABILITY STUDY

Intra-rater Reliability Study	Number of Measurements	Intraclass Correlation Coefficient ICC	95% Confidence Interval
Humac Norm Computerised Dynamometer	110 Repeated four days later	0.948	0.992 to 0.965

Recruitment and assessment of SSI cases and controls were conducted at the same time, independently of each other, with matching not performed until data collection

was completed. Seventy-three SSI cases and 91 controls were assessed and then matched for gender, hand dominance, physical activity level and age (within a bracket of three years). SSI cases reported symptoms being present between 4 weeks to 12 months. This resulted in 51 complete matches in each group. Within the SSI group, 31 dominant limbs were symptomatic and 20 non-dominant limbs were symptomatic. No significant differences in body mass index or physical activity was identified between the groups, with moderate activity level being the most prevalent in both groups (see Table 2). SPADI and VAS scores were significantly different (Table 2).

TABLE 2: COMPARISON OF SSI (CASES) AND CONTROL PARTICIPANTS

	SSI	CONTROL	P VALUE
	MEAN ± SD	MEAN ± SD	
	N = 51	N =51	
Age (years)	51.2 ± 5.7	50.8 ± 4.7	.074
BMI	28.1 ± 5.6	28.2 ± 4.6	.393
Gender			1.0
Male	28	28	
Female	23	23	
Dom			1.0
Right	45	45	
Left	6	6	
IPAQ			.282
Low	27%	30.2%	
Mod	42.9%	38.1%	

High	30.2%	31.7%	
VAS Rest	0.2 ± 0.8	0	.000
VAS Activity	5.8 ± 2.8	0	.000
SPADI	26.2 ± 17.9	0	.000

Abbreviations: BMI, body mass index; Dom, dominance; Asym, Asymptomatic; VAS, Visual Analogue Scale; SPADI, Shoulder Physical Activity Disability Index

SSI (Cases) versus Control Analysis

Dominant Shoulder

Significantly less con ER PT at 60⁰/second (p=0.025), ecc ER PT at 120⁰/second (p=0.015), ecc ER RPT at 120⁰/second (p=0.043) and ecc IR PT at 60⁰/second (p=0.013) and 120⁰/second (p=0.031) was identified in the dominant symptomatic SSI shoulder compared to the dominant control shoulder (table 3). While no other statistical differences were identified it was noted that all measures of the SSI dominant shoulder were lower than the dominant control shoulder.

Non-Dominant Shoulder

No significant difference in isokinetic strength was identified between the non-dominant SSI symptomatic shoulder and the non-dominant control shoulder. It is noted however that measurements for ER (PT and RPT, and both ratios in the SSI (cases) were higher in the control group whereas IR (PT and RPT) were slightly lower (table 3).

TABLE 3: ISOKINETIC TESTING FOR SYMPTOMATIC SHOULDER IN SSI GROUP (CASES) AND THE MATCHED SHOULDER IN CONTROL GROUP

OUTCOME MEASURE		SSI		CONTROL		95%CI	P VALUE
D (N=31) ND (N=20)		MEAN ± SD	(SEM)	MEAN ± SD	(SEM)		
PT ER Con							
60 ⁰ sec	D	12.7 ± 6.6	(1.2)	17.2 ± 8.4	(1.5)	-8.3 to -0.6	.025
	ND	16.0 ± 6.6	(1.5)	14.6 ± 6.8	(1.5)	-2.9 to 5.7	.511
120 ⁰ sec	D	10.7 ± 6.1	(0.1)	13.6 ± 7.8	(1.4)	-6.5 to 0.6	.103
	ND	11.7 ± 6.1	(1.4)	11.1 ± 6.0	(1.3)	-3.3 to 4.4	.775
PT ER Ecc							
60 ⁰ sec	D	21.6 ± 10.7	(1.9)	26.1 ± 11.0	(2.0)	-10.0 to 1.0	.109
	ND	25.5 ± 12.9	(3.0)	23.2 ± 9.1	(2.0)	-4.9 to 9.4	.529
120 ⁰ sec	D	21.3 ± 7.0	(1.2)	27.2 ± 11.1	(2.0)	-10.6 to -1.2	.015
	ND	29.1 ± 17.0	(3.8)	25.5 ± 9.4	(2.1)	-5.2 to 12.4	.411
Rel PT ER Con							
60 ⁰ sec	D	0.16 ± 0.08	(0.01)	0.20 ± 0.09	(0.02)	-0.08 to 0.00	.057
	ND	0.18 ± 0.07	(0.01)	0.17 ± 0.07	(0.01)	-0.03 to 0.05	.638
120 ⁰ sec	D	0.14 ± 0.07	(0.01)	0.16 ± 0.09	(0.02)	-0.07 to 0.01	.209
	ND	0.14 ± 0.07	(0.01)	0.14 ± 0.07	(0.01)	-0.04 to 0.04	.989
Rel PT ER Ecc							
60 ⁰ sec	D	0.28 ± 0.12	(0.02)	0.31 ± 0.10	(0.02)	-0.09 to 0.02	.242
	ND	0.30 ± 0.18	(0.04)	0.28 ± 0.08	(0.02)	-0.07 to 0.11	.607
120 ⁰ sec	D	0.27 ± 0.09	(0.02)	0.32 ± 0.10	(0.02)	-0.10 to -0.00	.043
	ND	0.35 ± 0.26	(0.06)	0.31 ± 0.09	(0.02)	-0.08 to 0.17	.477
PT IR Con							
60 ⁰ sec	D	31.9 ± 11.9	(2.1)	36.2 ± 13.6	(2.4)	-10.8 to 2.2	.195
	ND	33.1 ± 11.8	(2.6)	34.4 ± 14.3	(3.2)	-9.7 to 7.1	.755
120 ⁰ sec	D	31.1 ± 11.6	(2.1)	33.6 ± 14.1	(2.5)	-9.1 to 4.1	.452
	ND	32.2 ± 11.5	(2.6)	30.9 ± 15.1	(3.4)	-7.3 to 9.8	.770
PT IR Ecc							

60 ⁰ sec	D	39.9 ± 13.4	(2.4)	49.3 ± 15.7	(2.8)	-16.9 to -2.1	.013
	ND	45.8 ± 15.6	(3.5)	48.6 ± 17.7	(4.0)	-13.5 to 7.8	.593
120 ⁰ sec	D	41.8 ± 11.5	(2.1)	49.8 ± 16.6	(3.0)	-15.3 to -0.8	.031
	ND	47.2 ± 16.1	(3.6)	49.6 ± 18.3	(4.1)	-13.4 to 8.7	.669
Rel PT IR Con							
60 ⁰ sec	D	0.41 ± 0.16	(0.03)	0.44 ± 0.14	(0.02)	-0.10 to 0.06	.591
	ND	0.40 ± 0.17	(0.04)	0.43 ± 0.17	(0.04)	-0.13 to 0.08	.606
120 ⁰ sec	D	0.40 ± 0.14	(0.03)	0.40 ± 0.15	(0.03)	-0.08 to 0.07	.934
	ND	0.39 ± 0.11	(0.04)	0.38 ± 0.18	(0.04)	-0.10 to 0.11	.903
Rel PT IR Ecc							
60 ⁰ sec	D	0.52 ± 0.18	(0.03)	0.60 ± 0.15	(0.03)	-0.16 to 0.00	.058
	ND	0.54 ± 0.20	(0.05)	0.60 ± 0.19	(0.04)	-0.18 to 0.07	.412
120 ⁰ sec	D	0.54 ± 0.15	(0.03)	0.60 ± 0.16	(0.03)	-0.14 to 0.02	.137
	ND	0.55 ± 0.18	(0.04)	0.61 ± 0.19	(0.04)	-0.17 to 0.07	.374
Ratio ER Ecc/IR Con							
60 ⁰ sec	D	0.72 ± 0.32	(0.06)	0.74 ± 0.23	(0.04)	-0.17 to 0.11	.690
	ND	0.81 ± 0.44	(0.10)	0.71 ± 0.23	(0.05)	-0.12 to 0.32	.362
120 ⁰ sec	D	0.72 ± 0.22	(0.04)	0.91 ± 0.58	(0.10)	-0.41 to 0.03	.094
	ND	0.97 ± 0.58	(0.13)	0.94 ± 0.50	(0.11)	-0.31 to 0.38	.822
Ratio ER Con/IR Con							
60 ⁰ sec	D	0.41 ± 0.19	(0.03)	0.48 ± 0.17	(0.03)	-0.15 to 0.03	.162
	ND	0.52 ± 0.24	(0.05)	0.43 ± 0.14	(0.03)	-0.04 to 0.22	.163
120 ⁰ sec	D	0.34 ± 0.12	(0.02)	0.41 ± 0.15	(0.03)	-0.14 to 0.00	.058
	ND	0.38 ± 0.19	(0.04)	0.38 ± 0.16	(0.03)	-0.11 to 0.12	.952

Abbreviations: PT, Peak Torque; Rel PT, relative peak torque; ER, External Rotation; IR, Internal Rotation; Con, Concentric; Ecc, Eccentric; D, Dominant; ND, Non-Dominant

No significant differences were identified when the asymptomatic shoulder of the SSI (cases) (dominant = 20, non-dominant = 31) was compared with the matched shoulder of the control group (table 4).

TABLE 4: ISOKINETIC TESTING FOR ASYMPTOMATIC SHOULDERS IN SSI (CASES) AND MATCHED SHOULDERS IN THE CONTROL GROUP

OUTCOME MEASURE		SSI		CONTROL		95%CI	p value
D (n=20)	ND (n=31)	Mean ± SD (SEM)		Mean ± SD (SEM)			
PT ER Con							
60 ⁰ sec	D	17.0 ± 7.2	(1.6)	16.5 ± 6.7	(1.5)	-4.0 to 5.0	.822
	ND	12.3 ± 5.6	(1.0)	14.1 ± 6.5	(1.2)	-4.9 to 1.3	.257
120 ⁰ sec	D	13.5 ± 7.3	(1.6)	11.7 ± 6.7	(1.5)	-2.7 to 6.3	.422
	ND	10.9 ± 5.7	(1.0)	11.0 ± 6.2	(1.1)	-3.1 to 3.0	.983
PT ER Ecc							
60 ⁰ sec	D	26.8 ± 9.3	(2.1)	24.5 ± 9.0	(2.0)	-3.5 to 8.2	.420
	ND	21.6 ± 8.9	(1.6)	22.8 ± 8.7	(1.6)	-5.6 to 3.3	.605
120 ⁰ sec	D	28.8 ± 9.8	(2.2)	27.3 ± 12.4	(2.8)	-5.6 to 8.6	.673
	ND	22.1 ± 8.2	(1.5)	24.1 ± 9.3	(1.7)	-6.4 to 2.5	.381
Rel PT ER Con							
60 ⁰ sec	D	0.19 ± 0.07	(0.01)	0.20 ± 0.07	(0.02)	-0.05 to 0.04	.687
	ND	0.16 ± 0.07	(0.01)	0.17 ± 0.07	(0.01)	-0.05 to 0.03	.573
120 ⁰ sec	D	0.15 ± 0.07	(0.02)	0.14 ± 0.07	(0.02)	-0.03 to 0.06	.554
	ND	0.14 ± 0.07	(0.01)	0.13 ± 0.06	(0.01)	-0.02 to 0.04	.614
Rel PT ER Ecc							
60 ⁰ sec	D	0.31 ± 0.08	(0.02)	0.30 ± 0.09	(0.02)	-0.04 to 0.06	.709
	ND	0.28 ± 0.12	(0.02)	0.27 ± 0.09	(0.02)	-0.05 to 0.06	.879
120 ⁰ sec	D	0.33 ± 0.09	(0.02)	0.33 ± 0.12	(0.03)	-0.07 to 0.07	.930
	ND	0.28 ± 0.10	(0.02)	0.29 ± 0.09	(0.02)	-0.05 to 0.04	.813
PT IR Con							
60 ⁰ sec	D	34.9 ± 10.3	(2.3)	35.5 ± 18.3	(4.1)	-10.0 to 8.9	.907
	ND	34.3 ± 13.3	(2.4)	34.2 ± 13.2	(2.4)	-6.6 to 6.9	.962
120 ⁰ sec	D	30.9 ± 11.8	(2.6)	32.5 ± 15.8	(3.5)	-10.5 to 7.4	.727
	ND	32.5 ± 12.9	(2.3)	32.0 ± 14.4	(2.6)	-6.4 to 7.5	.452
PT IR Ecc							
60 ⁰ sec	D	49.5 ± 14.4	(3.2)	50.3 ± 21.6	(4.8)	-12.5 to 10.9	.891
	ND	43.3 ± 14.7	(2.6)	44.0 ± 15.0	(2.7)	-8.2 to 6.9	.858
120 ⁰ sec	D	50.6 ± 12.3	(2.8)	52.1 ± 18.3	(4.1)	-11.5 to 8.5	.763
	ND	44.7 ± 14.3	(2.6)	45.9 ± 15.3	(2.7)	-8.7 to 6.3	.752

Rel PT IR Con							
60 ⁰ sec	D	0.41 ± 0.12	(0.03)	0.44 ± 0.21	(0.05)	-0.14 to 0.08	.549
	ND	0.44 ± 0.17	(0.03)	0.41 ± 0.13	(0.02)	-0.04 to 0.11	.393
120 ⁰ sec	D	0.36 ± 0.14	(0.03)	0.40 ± 0.17	(0.04)	-0.14 to 0.06	.471
	ND	0.42 ± 0.16	(0.03)	0.38 ± 0.14	(0.03)	-0.04 to 0.11	.334
Rel PT IR Ecc							
60 ⁰ sec	D	0.57 ± 0.14	(0.03)	0.61 ± 0.22	(0.05)	-0.16 to 0.08	.479
	ND	0.56 ± 0.17	(0.03)	0.53 ± 0.14	(0.03)	-0.05 to 0.11	.517
120 ⁰ sec	D	0.59 ± 0.13	(0.03)	0.64 ± 0.19	(0.04)	-0.16 to 0.05	.306
	ND	0.57 ± 0.17	(0.03)	0.55 ± 0.14	(0.03)	-0.06 to 0.10	.586
Ratio ER Ecc/IR Con							
60 ⁰ sec	D	0.78 ± 0.20	(0.04)	0.81 ± 0.50	(0.11)	-0.27 to 0.22	.808
	ND	0.66 ± 0.28	(0.05)	0.69 ± 0.19	(0.03)	-0.15 to 0.10	.674
120 ⁰ sec	D	1.0 ± 0.39	(0.09)	0.92 ± 0.47	(0.10)	-0.20 to 0.35	.565
	ND	0.71 ± 0.25	(0.04)	0.84 ± 0.41	(0.07)	-0.30 to 0.05	.164
Ratio ER Con/IR Con							
60 ⁰ sec	D	0.48 ± 0.14	(0.03)	0.52 ± 0.21	(0.05)	-0.14 to 0.08	.585
	ND	0.36 ± 0.10	(0.02)	0.41 ± 0.14	(0.02)	-0.11 to 0.01	.089
120 ⁰ sec	D	0.42 ± 0.14	(0.03)	0.36 ± 0.17	(0.04)	-0.04 to 0.16	.233
	ND	0.33 ± 0.12	(0.02)	0.35 ± 0.13	(0.02)	-0.08 to 0.04	.532

Abbreviations: PT, Peak Torque; Rel PT, relative peak torque; ER, External Rotation; IR, Internal Rotation; Con, Concentric; Ecc, Eccentric; D, Dominant; ND, Non-Dominant

DISCUSSION

Rotator cuff weakness is reported to be associated with SSI (J.S. Lewis, 2009;

Michener et al., 2003) yet very few studies have investigated rotator cuff strength in

an SSI group and an asymptomatic group. The hypothesis that a significant

difference in muscle strength would be found in the painful shoulder in the SSI group

(cases) compared to the dominance matched shoulder in the control group has not

been clearly identified in this study. Only one concentric variable (ER PT at 60⁰

/second) was significantly different between the two groups. Yet concentric testing has been shown to be more reliable than eccentric testing when comparing an SSI group to an asymptomatic group (MacDermid et al., 2004). A concentric contraction produces less force than an eccentric contraction, thereby reducing the influence of pain on performance (Anderson, Bialocerkowski, & Bennell, 2006).

Only one previous study has compared concentric isokinetic ER and IR PT in an SSI symptomatic shoulder with a control group (Dulgeroglu et al., 2013). All PT values were found to be significantly lower for con IR and con ER, at 90⁰ sec and 180⁰ sec, in the SSI symptomatic shoulder compared to the dominant shoulder of the control group. However, of the 22 symptomatic shoulders assessed only 14 of these were actually the dominant shoulder (Dulgeroglu et al., 2013). The remaining eight shoulders were non-dominant, however these were compared to the dominant shoulders of the control group. This analytical and methodological anomaly, together with the relatively small sample size of the study, may explain why the findings of the current study differ from the results reported (Dulgeroglu et al., 2013)

Differences in eccentric strength in this study were only present when the dominant shoulder was the effected shoulder in the SSI group. Significantly less ecc ER PT at 120⁰/second, ecc ER RPT at 120⁰/second and ecc IR PT at 60⁰/second and 120⁰/second was found when compared to the matched control shoulder.

When the non-dominant shoulder was the affected shoulder in the SSI (cases) no significant differences were identified compared to the matched non-dominant shoulder in the control group. No previous studies have been identified which have directly compared these variables in an SSI group and a control group.

Mean values for all measurements of the dominant shoulder in the SSI (cases) were consistently lower compared to the matched dominant shoulder in the control group.

However, when the non-dominant shoulder was the affected shoulder in the SSI (cases) the values were very similar or slightly higher compared to the control group. It appears that changes in strength in SSI are related to the dominance of the SSI symptomatic shoulder, which may have implications for strengthening regimes.

Isokinetic testing in a SSI group and an asymptomatic group using a similar age group, tested in the seated position, with the shoulder positioned in the scapular plane, has been reported in five previous studies (Dulgeroglu et al., 2013; Erol et al., 2008; Leroux et al., 1994; MacDermid et al., 2004; Tyler et al., 2005).

One study reported within group differences of an SSI group compared to within group differences of an asymptomatic group (Erol et al., 2008). Only right hand dominant participants were recruited to both groups and matched for age, sex, height and body weight, with concentric testing performed at 60⁰ sec. No within group difference between dominant and non-dominant limbs in the SSI group was identified whereas a significant difference was found in the asymptomatic group. The similarity in findings with this study, albeit a small sample size (13 SSI, 25 control), likely reflects the same methodology of matched limb dominance, age and gender.

Other studies which used the same isokinetic testing position to compare an SSI group with an asymptomatic group did not report dominance of the recruited participants but then analysed within group differences for the (1) painful and non-painful shoulders in those with SSI and (2) dominant versus non-dominant shoulder in an asymptomatic group and then (3) compared the values from these two analyses (Leroux et al., 1994; MacDermid et al., 2004; Tyler et al., 2005). This statistical analysis differs from the analysis in this study. The findings of these previous studies are difficult to compare to the outcomes of this study as limb

dominance and the presence of pain will both have an effect on isokinetic performance.

Limitations of this study include the availability of only one assessor to perform all isokinetic testing, however extraction of computer generated data was checked by an independent assessor. Another limitation was the participants not being familiar with the use of the isokinetic dynamometer which is in common with other isokinetic studies. Although instructions were clear before commencing the trial reminders to apply maximum effort throughout and which direction to apply resistance were sometimes needed for both those in the SSI group and the control group. However, this was true for both cases and controls so the measurement bias is likely to be non-differential. The effect of pain was minimized by the position and range chosen for testing. Selection bias (volunteer bias) may be present due to a snowballing effect recruitment strategy. This study only included participants aged 40 to 60 years. While the primary age of SSI, these findings should only be applied to this age group. A strength of this study is the matching of cases and controls on age, gender, hand dominance and physical activity levels.

CONCLUSION

This study is the first to compare isokinetic rotator cuff testing at 60⁰ and 120⁰ per second through a total range of 60⁰ in 40 to 60 year olds experiencing SSI and a control group matched for age, gender, hand dominance and physical activity levels.

Differences in muscle strength were not clearly identified between the SSI cases and control group with significant strength differences only found when the dominant SSI shoulder was symptomatic (con ER PT at 60⁰/second, ecc ER at 120⁰/second, ecc ER RPT at 120⁰/second and ecc IR PT at 60⁰/second and 120⁰/second). No strength differences were evident when comparing the non-dominant symptomatic

SSI shoulder and the non-dominant control shoulder indicating strength in SSI may be related to dominance, which may have implications for strengthening regimes.

APPENDIX A

Methodological Detail

Isokinetic Assessment

Isokinetic testing was performed using a Humac Norm Computerised Dynamometer ((CSMI), 2006). Calibration was completed prior to testing taking place. The asymptomatic group were randomly allocated by drawing a piece of paper from a box to determine the arm to be tested first. The asymptomatic limb was consistently tested first in the SIS group, with this familiarisation encouraging maximal effort when testing the symptomatic limb.

The participant was seated in the standardised position ensuring the seat position allowed the testing arm to be at 45 degrees abduction in the scapular plane. The set up was consistent with those provided in the Humac Norm System User's Guide, page 5-34 ((CSMI), 2006). The chair was rotated to 35⁰, dyna tilt was 45⁰ and dyna rotation was 5⁰. A heat moulded wrist splint was attached before the arm was positioned and strapped into place. Initial recruits displayed visible flexion and extension occurring at the wrist during testing. A small study was then conducted, using asymptomatic young participants, performing the same protocol with the splint in situ and without and no significant difference in peak torque values was found at either speed. However, it was decided to use the heat moulded splint for all participants to standardise the wrist joint position. The zero rotation position was established using a spirit level resting on the fixed arm attachment of the machine.

Standardised instructions were given by the examiner explaining which direction the movement was to occur, to provide maximum effort, and keep the pressure throughout the entire movement. 3 practice reps were allowed before each test. The examiner advised not to provide maximum effort in the practice reps but just get used to the machine. One minute rest was provided between practice and trial. Five trials were done in each direction. These were reciprocal concentric/eccentric external rotation and concentric/eccentric internal rotation at 60 degrees per second. One minute rest was then given followed by reciprocal concentric/eccentric external rotation and concentric/eccentric internal rotation at 120 degrees per second. All tests were completed on one arm before adjusting the seat set –up to allow testing with the other arm. All 5 repetitions at each speed were included in the analysis.



FIGURE 1. Humac Norm Set Up

- (CSMI), C. S. M. I. (2006). *Humac Norm Testing & Rehabilitation System Users Guide Model 770*. Stoughton, MA: Computer Sports Medicine Inc.
- Altman, D. G. (1991). *Practical statistics for medical research*: Chapman & Hall.
- Anderson, V. B., Bialocerowski, A. E., & Bennell, K. L. (2006). Test-retest reliability of glenohumeral internal and external rotation strength in chronic rotator cuff pathology. *Physical Therapy In Sport*, 7, 115-121. doi: 10.1016/j.ptsp.2006.04.002
- Bigliani, L. U., & Levine, W. N. (1997). Subacromial Impingement Syndrome. *Journal of Bone and Joint Surgery*, 79(12), 1854-1868.
- Bygott, I. L., McMeeken, J., Carroll, S., & Story, I. (2001). Gravity correction in trunk dynamometry: Is it reliable? *Isokinetics and Exercise Science*, 9, 1-9.
- Craig, C. L., Marshall, A. L., Sjostrom, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., . . . Oja, P. (2003). International physical activity questionnaire: 12 country reliability and validity. *Medicine and Science in Sports and Exercise*, 35, 1381-1395. doi: 10.1249/01.MSS.0000078924.61453.FB
- Dalton, S. E. (1989). Clinical examination of the painful shoulder. *Ballieres Clinical Rheumatology*, 3(3), 453-474.
- Dark, A., Ginn, K. A., & Halaki, M. (2007). Shoulder Muscle Recruitment Patterns During Commonly Used Rotator Cuff Exercises: An Electromyographic Study. *Physical Therapy*, 87(8), 1039-1046.
- David, G., Magarey, M. E., Jones, M. A., Dvir, Z., Turker, K. S., & Sharpe, M. (2000). EMG and strength correlates of selected shoulder muscles during rotations of the glenohumeral joint. *Clinical Biomechanics*, 15, 95 - 102.
- Dogu, B., Sahin, F., Ozmaden, A., Yilmaz, F., & Kuran, B. (2013). Which questionnaire is more effective for follow-up diagnosed subacromial impingement syndrome? A comparison of the responsiveness of SDQ, SPADI and WORC index. *Journal of Back and Musculoskeletal Rehabilitation*, 26, 1-7. doi: 10.3233/BMR-2012-0342
- Dube, J. A., & Mercier, C. (2011). Effect of pain and pain expectation on primary motor cortex. *Clinical Neurophysiology*, 122, 2318-2323. doi: 10.1016/j.clinph.2011.03.026
- Dulgeroglu, D., Kirbiyik, E., Ersoz, M., & Ozel, S. (2013). Evaluation of shoulder rotational strength in patients with subacromial impingement syndrome using a computerized isokinetic dynamometer. *Journal of musculoskeletal pain*, 21(1), 23-30. doi: 10.3109/10582452.2013.763393
- Erol, O., Ozcakar, L., & Celiker, R. (2008). Shoulder rotator cuff strength in patients with stage I-II subacromial impingement: Relationship to pain, disability and quality of life. *Journal of shoulder and elbow surgery*, 17(6), 893-897. doi: 10.1016/j.jse.2008.05.043
- Hanchard, N., Cummins, J., & Jeffries, C. (2004). Evidence-based clinical guidelines for the diagnosis, assessment and physiotherapy management of shoulder impingement syndrome. Retrieved from
- Hanchard, N. C. A., & Handoll, H. H. G. (2008). Physical tests for shoulder impingements and local lesions of bursa, tendon or labrum that may accompany impingement. *Cochrane Database of Systematic Reviews*(4). doi: 10.1002/14651858.CD007427
- Hawkins, R. J., & Kennedy, J. C. (1980). Impingement syndrome in athletes. *American Journal of Sports Medicine*, 8(3), 151-158. doi: 10.1177/036354658000800302
- Holmgren, T., Bjornsson Hallgren, H., Oberg, B., Adolfsson, L., & Johansson, K. (2012). Effect of specific exercise strategy on need for surgery in patients with subacromial impingement

- syndrome: Randomized Control Study. *British Medical Journal*(BMJ 2012;344:e787), 9. doi: 10.1136/bmj.e787
- Jensen, M. P., Karoly, P., & Braver, S. (1986). The measurement of clinical pain intensity: a comparison of six methods. *Pain*, 27, 117-126.
- Kessel, L., & Watson, M. (1977). The painful arc syndrome. Clinical classification as a guide to management. *The Journal of Bone and Joint Surgery*, 59-B(2), 166-172.
- Kuhlman, J. R., Iannotti, J. P., Kelly, M. J., Riegler, F. X., Gevaert, M. L., & Ergin, T. M. (1992). Isokinetic and isometric measurement of strength of external rotation and abduction of the shoulder. *Journal of Bone and Joint Surgery American*, 74, 1320-1333.
- Kuhn, J. E. (2009). Exercise in the treatment of rotator cuff impingement: A systematic review and a synthesized evidence-based rehabilitation protocol. *Journal of shoulder and elbow surgery*, 18, 138-160.
- Land, H., & Gordon, S. (2011). What is normal isokinetic shoulder strength or strength ratios? A systematic review. *Isokinetics and Exercise Science*, 19, 231-241. doi: 10.3233/IES-2011-0427
- Leroux, J.-L., Codine, P., Thomas, E., Pocholle, M., Mailhe, D., & Blotman, F. (1994). Isokinetic evaluation of rotational strength in normal shoulders and shoulders with impingement syndrome. *Clinical Orthopaedics and Related Research*, 108-115.
- Lewis, J., & Ginn, K. (2015). Rotator cuff tendinopathy and subacromial pain syndrome. In G. Jull, A. Moore, D. Falla, J. Lewis, C. McCarthy, & M. Sterlin (Eds.), *Grieve's Modern Musculoskeletal Physiotherapy* (Fourth ed., pp. 563-568): Elsevier.
- Lewis, J. S. (2009). Rotator cuff tendinopathy (Vol. 43, pp. 236-241).
- Lewis, J. S. (2009). Rotator Cuff Tendinopathy/Subacromial Impingement Syndrome. Is it time for a new method of assessment? *British Journal of Sports Medicine*, 43, 259-264.
- Lewis, J. S., Green, A. S., & Dekel, S. (2001). The aetiology of subacromial impingement syndrome. *Physiotherapy*, 87(9), 458-469.
- Ludewig, P. M., & Cook, T. M. (2000). Alterations in shoulder kinematics and associated muscle activity in people with symptoms of shoulder impingement. *Physical Therapy*, 276-291.
- MacDermid, J. C., Ramos, J., Drosdowech, D., Faber, K. J., & Patterson, S. (2004). The impact of rotator cuff pathology on isometric and isokinetic strength, function and quality of life. *Journal of shoulder and elbow surgery*, 13(6), 593-598. doi: 10.1016/j.jse.2004.03.009
- Michener, L. A., McClure, P. W., & Karduna, A. R. (2003). Anatomical and biomechanical mechanisms of subacromial impingement syndrome. *Clinical Biomechanics*, 18, 369-379.
- Michener, L. A., Walsworth, M. K., Doukas, W. C., & Murphy, K. P. (2009). Reliability and diagnostic accuracy of 5 physical examination tests and combination of tests for subacromial impingement. *Archives of Physical Medicine and Rehabilitation*, 90, 1898-1903. doi: 10.1016/j.apmr.2009.05.015
- Neer, C. S. I. (1983). Impingement Lesions. *Clinical Orthopaedics and Related Research*, 173(March), 70-77.
- Ngomo, S., Mercier, C., Bouyer, L. J., Sacoie, A., & Roy, J.-S. (2015). Alterations in central motor representation increase over time in individuals with rotator cuff tendinopathy. *Clinical Neurophysiology*, 126, 365-371. doi: 10.1016/j.clinph.2014.05.035
- Ostor, A. J. K., Richards, C. A., Prevost, A. T., Speed, C. A., & Hazleman, B. L. (2005). Diagnosis and relation to general health of shoulder disorders presenting to primary care. *Rheumatology*, 44, 800-805. doi: 10.1093/rheumatology/keh598
- Park, H. B., Yokota, A., Gill, H. S., Rassi, G. E., & McFarland, E. G. (2005). Diagnostic accuracy of clinical tests for the different degrees of subacromial impingement syndrome. *Journal of Bone and Joint Surgery*, 87-A(7), 1446-1455.
- Reddy, A. S., Mohr, K. R., Pink, M. M., & Jobe, F. W. (2000). Electromyographic analysis of the deltoid and rotator cuff muscles in persons with subacromial impingement. *Journal of shoulder and elbow surgery*, 9, 519-523.

- Reinold, M. M., Macrina, L. C., Wilk, K. E., Fleisig, G. S., Dun, S., Barrentine, S. W., . . . Andrews, J. R. (2007). Electromyographic analysis of the supraspinatus and deltoid muscles during 3 common rehabilitation exercises. *Journal of Athletic Training, 42*(4), 464-469.
- Reinold, M. M., Wilk, K. E., Fleisig, G. S., Zheng, N., Barrentine, S. W., Chmielewski, T., . . . Andrews, J. R. (2004). Electromyographic analysis of the rotator cuff and deltoid musculature during common shoulder external rotation exercises. *Journal of Orthopaedic & Sports Physical Therapy, 34*(7), 385-394.
- Roach, K. E., Budiman-Mak, E., Songsiridej, N., & Lertratanakul, Y. (1991). Development of a shoulder pain and disability index. *Arthritis & Rheumatism, 4*(4), 143-149.
- Roy, J.-S., Macdermid, J. C., Boyd, K. U., Faber, K. J., Drosdowech, D., & Athwal, G. S. (2009). Rotational strength, range of motion, and function in people with unaffected shoulders from various stages of life. *Sports Medicine, Arthroscopy, Rehabilitation, Therapy and Technology, 1*(4). doi: 10.1186/1758-2555-1-4
- Roy, J.-S., Moffet, H., & McFadyen, B. J. (2008). Upper limb motor strategies in persons with and without shoulder impingement syndrome across different speeds of movement. *Clinical Biomechanics, 23*, 1227-1236. doi: 10.1016/j.clinbiomech.2008.07.009
- Tate, A. R., McClure, P. W., Young, I. A., Salvatori, R., & Michener, L. A. (2010). Comprehensive impairment-based exercise and manual therapy intervention for patients with subacromial impingement syndrome: A case series. *Journal of Orthopaedic & Sports Physical Therapy, 40*(8), 474-493. doi: 10.2519/jospt.2010.3223
- Tyler, T. F., Nahow, R. C., Nicholas, S. J., & McHugh, M. P. (2005). Quantifying shoulder rotation weakness in patients with shoulder impingement. *Journal of shoulder and elbow surgery, 14*(6), 570-574.
- van der Windt, A. W. M., Koes, B. W., de Jong, B. A., & Bouter, L. M. (1995). Shoulder disorders in general practice: incidence, patient characteristics, and management. *Annals of the Rheumatic Diseases, 54*, 959-964. doi: 10.1136/ard.54.12.959
- van Meeteren, J., Roebroek, M. E., Selles, R. W., Stijnen, T., & Stem, H. J. (2004). Concentric isokinetic dynamometry of the shoulder: Which parameters discriminate between healthy subjects and patients with shoulder disorders? *Isokinetics and Exercise Science, 12*, 239-246.